

POLYGONS THAT GENERATE AN EDGE TESSELLATION OF THE PLANE

ANDREW HALL, MATTHEW KIRBY, JOSHUA YORK, AND RON UMBLE

ABSTRACT. An *edge tessellation of the plane* is generated by the reflection of some polygon in its edges. We prove that a polygon generating an edge tessellation is exactly one of the following: An equilateral, 30-right, isosceles right, or 120-isosceles triangle, a rectangle, a 60-rhombus, a 60-90-120-kite, or a regular hexagon.

The results in this paper were discovered by Millersville University students Andrew Hall, Matthew Kirby, and Joshua York during an undergraduate research seminar directed by the fourth author in the spring of 2009. The goal of the seminar was to identify those polygons that generate a tessellation of the plane when reflected in their edges. Such tessellations, referred to here as *edge tessellations*, are examples of Laves tilings [3]. This problem is motivated by recent work by A. Baxter and R. Umble who use an edge tessellation to find, classify, and count classes of periodic orbits on equilateral triangles [1]. Indeed, a polygon to which their classification techniques can be applied necessarily generates an edge tessellation.

Consider a polygon G that generates an edge tessellation. Observe that the measures of the interior angles of G lie in the set $\{30^\circ, 45^\circ, 60^\circ, 90^\circ, 120^\circ\}$. Let θ be the measure of the interior angle at some vertex V of G , and note that $\theta < 180^\circ$. If G' is the reflection of G in an edge containing V , the interior angle of G' at V has measure θ , and inductively, every interior angle at V has measure θ . Thus V is a point of n -fold symmetry since successively reflecting in the edges of G that meet at V is a rotational symmetry. But $n = 2, 3, 4, 6$ by Crystallographic Restriction [4]. Consequently, if G' is the image of G under a rotational symmetry, the minimum rotation angle is θ and V is shared by 3, 4, or 6 copies of G . Note that when $n = 3$, exactly three copies of G share vertex V and G is symmetric with respect to its angle bisector at V . Otherwise, G and G' rotate together as a unit, in which case the minimum rotation angle is 2θ and V is shared by 4, 6, 8, or 12 copies of G . The conclusion now follows.

Next, let e be the number of edges of G ; then the interior angle sum $S = 180(e - 2)$. On the other hand, $S \leq 120e$ since the interior angles measure at most 120° . But $180(e - 2) \leq 120e$ implies $e \leq 6$ and G has at most 6 edges. Furthermore, it is a well-known fact that there are no regular tilings of the plane by pentagons [4]. Therefore G is a triangle, a quadrilateral, or a hexagon.

The fact that G has 3, 4, or 6 edges, has interior angles measuring 30° , 45° , 60° , 90° , or 120° , and is symmetric with respect to its angle bisectors at interior angles of 120° proves:

Date: July 10, 2009.

Key words and phrases. Tessellation, symmetry.

Proposition 1. *A polygon with an interior angle of 120° that generates an edge tessellation is either a 120-isosceles triangle, a 60-rhombus, a 60-90-120-kite, or a regular hexagon.*




Figure 1. Edge tessellations with an interior angle of 120° .

In [2], A. Beardon proved that a polygon whose interior angles measure $\frac{\pi}{n}$ with $n \in \mathbb{Z}$ generates an edge tessellation. Since angles measuring 30° , 45° , 60° , and 90° have this property, our problem reduces to determining all triangles and quadrilaterals whose interior angles have these particular measures. Let a , b , c , and d denote the number of 30° , 45° , 60° , and 90° interior angles of G , respectively, then each non-negative solution of the following system of equations uniquely determines a family of “similar” polygons with the desired property:

$$(0.1) \quad \begin{array}{rcccccccl} 30a & + & 45b & + & 60c & + & 90d & = & 180(e-2) \\ a & + & b & + & c & + & d & = & e. \end{array}$$

Our discussion above revealed that the only obtuse triangle generating an edge tessellations is 120-isosceles. To determine the non-obtuse triangles with this property, set $e = 3$ in system (0.1) and obtain the general solution

$$(0.2) \quad (a, b, c, d) = s(1, -2, 1, 0) + t(3, -4, 0, 1) + (-3, 6, 0, 0).$$

Since t is the number of 90° angles, $0 \leq t \leq 1$. When $t = 0$, solution (0.2) reduces to $(a, b, c, d) = s(1, -2, 1, 0) + (-3, 6, 0, 0)$. Then clearly, $s = 3$ and we obtain $(a, b, c, d) = (0, 0, 3, 0)$, confirming the fact that equilateral triangles generate edge tessellations. When $t = 1$, solution (0.2) reduces to $(a, b, c, d) = s(1, -2, 1, 0) +$

$(0, 2, 0, 1)$. Since s is the number of 60° angles, $0 \leq s \leq 1$ and

$$(a, b, c, d) = \begin{cases} (0, 2, 0, 1), & s = 0, \\ (1, 0, 1, 1), & s = 1, \end{cases}$$

indicating that 30-right and isosceles-right triangles generate edge tessellations (see Figure 2). Having identified all triangles that generate edge tessellations, we have proved:

Proposition 2. *A triangle that generates an edge tessellation is equilateral, 30-right, isosceles-right, or 120-isosceles.*

Quadrilaterals that generate edge tessellations are found in a similar way. The general solution of system (0.1) with $e = 4$ is

$$(0.3) \quad (a, b, c, d) = s(1, -2, 1, 0) + t(3, -4, 0, 1) + (-12, 16, 0, 0).$$

Note that $0 \leq t \leq 4$ since a quadrilateral has at most four interior angles measuring 90° . But $a = s + 3t - 12 \geq 0$ and $b = -2s - 4t + 16 \geq 0$ imply $t \geq 4$. Consequently $t = 4$, $s = 0$, and $(a, b, c, d) = (0, 0, 0, 4)$, indicating that rectangles are the only additional quadrilaterals that generate edge tessellations. Thus we have proved:

Proposition 3. *A quadrilateral that generates an edge tessellation is a rectangle, a 60-rhombus, or a 60-90-120 kite.*




Figure 2. Edge tessellations with interior angles at most 90° .

We conclude by collecting our results in the previous three propositions:

Theorem 1. *A polygon generating an edge tessellation is exactly one of the following: An equilateral, 30-right, isosceles-right, or 120-isosceles triangle, a rectangle, a 60-rhombus, a 60-90-120-kite, or a regular hexagon.*

REFERENCES

- [1] A. Baxter and R. Umble, Periodic Orbits for Billiards on an Equilateral Triangle, *Am. Math. Monthly*, **115** (6) (2008) 479-491.
- [2] A. F. Beardon, *The Geometry of Discrete Groups*, GTM 91, Springer-Verlag, New York, 1995.
- [3] B. Grünbaum and G. Shephard, *Tilings and Patterns*, W. H. Freeman and Company, New York, 1986.
- [4] G. E. Martin, *Transformation Geometry: An Introduction to Symmetry*, Springer-Verlag, New York, 1882.

DEPARTMENT OF MATHEMATICS, MILLERSVILLE UNIVERSITY OF PENNSYLVANIA, MILLERSVILLE,
PA. 17551

E-mail address: `ron.umble@millersville.edu`